

Algorithms for Sensor-Based Robotics: Potential Functions

Computer Science 336

<http://www.cs.jhu.edu/~hager/Teaching/cs336>


Professor Hager

<http://www.cs.jhu.edu/~hager>

CS 336/436, GD Hager with slides from Ji Yeong Lee, Howie Choset and Z. Dodds



The Basic Idea

- A really simple idea:
 - Suppose the goal is a point $g \in \mathbb{R}^2$
 - Suppose the robot is a point $r \in \mathbb{R}^2$
 - Think of a “spring” drawing the robot toward the goal and away from obstacles:
 - Can also think of like and opposite charges



Another Idea

- Think of the goal as the bottom of a bowl
- The robot is at the rim of the bowl
- What will happen?



CS 336/436, GD Hager with slides from Ji Yeong Lee, Howie Choset and Z. Dodds

The General Idea

- Both the bowl and the spring analogies are ways of storing potential *energy*
- The robot moves to a lower energy configuration
- A *potential function* is a function $U : \mathfrak{R}^m \rightarrow \mathfrak{R}$
- Energy is minimized by following the negative *gradient* of the potential energy function:

$$\nabla U(q) = DU(q)^T = \left[\frac{\partial U}{\partial q_1}(q), \dots, \frac{\partial U}{\partial q_m}(q) \right]^T$$

- We can now think of a *vector field* over the space of all q 's ...
 - at every point in time, the robot looks at the vector at the point and goes in that direction

CS 336/436, GD Hager with slides from Ji Yeong Lee, Howie Choset and Z. Dodds

Attractive/Repulsive Potential Field

$$U(q) = U_{\text{att}}(q) + U_{\text{rep}}(q)$$

- U_{att} is the “attractive” potential --- move to the goal
- U_{rep} is the “repulsive” potential --- avoid obstacles

CS 336/436, GD Hager with slides from Ji Yeong Lee, Howie Choset and Z. Dodds

Artificial Potential Field Methods: Attractive Potential


Conical Potential

$$U(q) = \zeta d(q, q_{\text{goal}}).$$

$$\nabla U(q) = \frac{\zeta}{d(q, q_{\text{goal}})}(q - q_{\text{goal}}).$$

Quadratic Potential

$$U_{\text{att}}(q) = \frac{1}{2}\zeta d^2(q, q_{\text{goal}}),$$



$$\begin{aligned} F_{\text{att}}(q) &= \nabla U_{\text{att}}(q) = \nabla \left(\frac{1}{2}\zeta d^2(q, q_{\text{goal}}) \right), \\ &= \frac{1}{2}\zeta \nabla d^2(q, q_{\text{goal}}), \\ &= \zeta(q - q_{\text{goal}}), \end{aligned}$$


CS 336/436, GD Hager with slides from Ji Yeong Lee, Howie Choset and Z. Dodds

Artificial Potential Field Methods: Attractive Potential

Combined Potential

$$U_{\text{att}}(q) = \begin{cases} \frac{1}{2}\zeta d^2(q, q_{\text{goal}}), & d(q, q_{\text{goal}}) \leq d_{\text{goal}}^*, \\ d_{\text{goal}}^* \zeta d(q, q_{\text{goal}}) - \frac{1}{2}\zeta (d_{\text{goal}}^*)^2, & d(q, q_{\text{goal}}) > d_{\text{goal}}^*. \end{cases}$$

$$\nabla U_{\text{att}}(q) = \begin{cases} \zeta(q - q_{\text{goal}}), & d(q, q_{\text{goal}}) \leq d_{\text{goal}}^*, \\ \frac{d_{\text{goal}}^* \zeta (q - q_{\text{goal}})}{d(q, q_{\text{goal}})}, & d(q, q_{\text{goal}}) > d_{\text{goal}}^*. \end{cases}$$




In some cases, it may be desirable to have distance functions that grow more slowly to avoid huge velocities far from the goal

one idea is to use the quadratic potential near the goal ($< d^*$) and the conic farther away
One minor issue: what?

CS 336/436, GD Hager with slides from Ji Yeong Lee, Howie Choset and Z. Dodds

The Repulsive Potential




$$U_{\text{rep}}(q) = \begin{cases} \frac{1}{2}\eta \left(\frac{1}{D(q)} - \frac{1}{Q^*} \right)^2, & D(q) \leq Q^*, \\ 0, & D(q) > Q^*, \end{cases}$$

whose gradient is

$$\nabla U_{\text{rep}}(q) = \begin{cases} \eta \left(\frac{1}{Q^*} - \frac{1}{D(q)} \right) \frac{1}{D^2(q)} \nabla D(q), & D(q) \leq Q^*, \\ 0, & D(q) > Q^*, \end{cases}$$

Repulsive Potential




CS 336/436, GD Hager with slides from Ji Yeong Lee, Howie Choset and Z. Dodds

Total Potential Function


$$U(q) = U_{\text{att}}(q) + U_{\text{rep}}(q)$$

$$F(q) = -\nabla U(q)$$



CS 336/436, GD Hager with slides from Ji Yeong Lee, Howie Choset and Z. Dodds

Potential Fields




CS 336/436, GD Hager with slides from Ji Yeong Lee, Howie Choset and Z. Dodds

Gradient Descent

- A simple way to get to the bottom of a potential

$$\dot{c}(t) = -\nabla U(c(t)).$$

- A *critical point* is a point x s.t. q^* where $\nabla U(q^*) = 0$.
 - Equation is stationary at a critical point
 - Max, min, saddle
 - Stability?



CS 336/436, GD Hager with slides from Ji Yeong Lee, Howie Choset and Z. Dodds

The Hessian


- For a 1-d function, how do we know we are at a unique minimum (or maximum)?
- The Hessian is the $m \times m$ matrix of second derivatives
- If the Hessian is nonsingular ($\text{Det}(H) \neq 0$), the critical point is a unique point
 - if H is positive definite ($x^t H x > 0$), a minimum
 - if H is negative definite, a maximum
 - if H is indefinite, a saddle point
- A function is **Morse** if its critical points are all isolated points (Hessian is everywhere non-singular).

CS 336/436, GD Hager with slides from Ji Yeong Lee, Howie Choset and Z. Dodds


Gradient Descent

Gradient Descent:


- $q(0) = q_{\text{start}}$
- $i = 0$
- while $\nabla U(q(i)) \neq 0$ do
 - $q(i+1) = q(i) - \alpha(i) \nabla U(q(i))$
 - $i = i+1$




(a)



(b)



(c)




(d)

CS 336/436, GD Hager with slides from Ji Yeong Lee, Howie Choset and Z. Dodds


Gradient Descent

Gradient Descent:


- $q(0) = q_{\text{start}}$
- $i = 0$
- while $\|\nabla U(q(i))\| > \epsilon$ do
 - $q(i+1) = q(i) - \alpha(i) \nabla U(q(i))$
 - $i = i+1$




(a)



(b)




(c)



(d)

CS 336/436, GD Hager with slides from Ji Yeong Lee, Howie Choset and Z. Dodds

Numerically “Smoother” Path




CS 336/436, GD Hager with slides from Ji Yeong Lee, Howie Choset and Z. Dodds

Single Object Distance

$$d_i(q) = \min_{c \in \mathcal{QO}_i} d(q, c) \quad \nabla d_i(q) = \frac{q - c}{d(q, c)}$$


$$U_{\text{rep}_i}(q) = \begin{cases} \frac{1}{2} \eta \left(\frac{1}{d_i(q)} - \frac{1}{Q_i^*} \right)^2, & \text{if } d_i(q) \leq Q_i^* \\ 0, & \text{if } d_i(q) > Q_i^* \end{cases}$$



$$U_{\text{rep}}(q) = \sum_{i=1}^n U_{\text{rep}_i}(q)$$

CS 336/436, GD Hager with slides from Ji Yeong Lee, Howie Choset and Z. Dodds

Compute Distance: Sensor Information



CS 336/436, GD Hager with slides from Ji Yeong Lee, Howie Choset and Z. Dodds

Computing Distance: Use a Grid

- use a discrete version of space and work from there
 - The Brushfire algorithm is one way to do this
 - need to define a grid on space
 - need to define connectivity (4/8)
 - obstacles start with a 1 in grid; free space is zero

n1	n2	n3
n4	n5	n6
n7	n8	n9

4

n1	n2	n3
n4	n5	n6
n7	n8	n9

8


CS 336/436, GD Hager with slides from Ji Yeong Lee, Howie Choset and Z. Dodds

Brushfire Algorithm

- Initially: create a queue L of pixels on the boundary of all obstacles
- While $L \neq \emptyset$
 - pop the top element t of L
 - if $d(t) = 0$,
 - set $d(t)$ to $1 + \min_{t' \in N(t), d(t') \neq 0} d(t')$
 - Add all $t' \in N(t)$ with $d(t') = 0$ to L (at the end)
- The result is a distance map d where each cell holds the minimum distance to an obstacle.
- The gradient of distance is easily found by taking differences with all neighboring cells.

CS 336/436, GD Hager with slides from Ji Yeong Lee, Howie Choset and Z. Dodds


Brushfire example



CS 336/436, GD Hager with slides from Ji Yeong Lee, Howie Choset and Z. Dodds

Potential Functions Question

- How do we know that we have only a single (global) minimum



- We have two choices:
 - not guaranteed to be a global minimum: do something other than gradient descent (what?)
 - make sure only one global minimum (a navigation function, which we'll see later).

The Wave-front Planner

- Apply the brushfire algorithm starting from the goal
- Label the goal pixel 2 and add all zero neighbors to L
 - While $L \neq \emptyset$
 - pop the top element of L, t
 - set $d(t)$ to $1 + \min_{t' \in N(t), d(t') > 1} d(t')$
 - Add all $t' \in N(t)$ with $d(t') = 0$ to L (at the end)
- The result is now a distance for every cell
 - gradient descent is again a matter of moving to the neighbor with the lowest distance value

CS 336/436, GD Hager with slides from Ji Yeong Lee, Howie Choset and Z. Dodds

The Wavefront Planner: Setup

7	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
6	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
5	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
4	0	0	0	0	1	1	1	1	1	1	1	1	0	0	0	
3	0	0	0	0	1	1	1	1	1	1	1	1	0	0	0	
2	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	2	
	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15

CS 336/436, GD Hager with slides from Ji Yeong Lee, Howie Choset and Z. Dodds

The Wavefront in Action (Part 1)

- Starting with the goal, set all adjacent cells with "0" to the current cell + 1
 - 4-Point Connectivity or 8-Point Connectivity?
 - Your Choice. We'll use 8-Point Connectivity in our example

7	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
6	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
5	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
4	0	0	0	0	1	1	1	1	1	1	1	1	0	0	0	0	0
3	0	0	0	0	1	1	1	1	1	1	1	1	0	0	0	0	0
2	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	3	3	0
0	0	0	0	0	0	0	0	0	0	0	0	0	0	3	2	0	0
	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	

CS 336/436, GD Hager with slides from Ji Yeong Lee, Howie Choset and Z. Dodds

The Wavefront in Action (Part 2)

- Now repeat with the modified cells
 - This will be repeated until no 0's are adjacent to cells with values ≥ 2
 - 0's will only remain when regions are unreachable

7	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
6	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
5	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
4	0	0	0	0	1	1	1	1	1	1	1	1	0	0	0	0	0
3	0	0	0	0	1	1	1	1	1	1	1	1	0	0	0	0	0
2	0	0	0	0	0	0	0	0	0	0	0	0	0	4	4	4	0
1	0	0	0	0	0	0	0	0	0	0	0	0	0	4	3	3	0
0	0	0	0	0	0	0	0	0	0	0	0	0	0	4	3	2	0
	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	

CS 336/436, GD Hager with slides from Ji Yeong Lee, Howie Choset and Z. Dodds

The Wavefront in Action (Part 3)

- Repeat again...

7	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
6	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
5	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
4	0	0	0	0	1	1	1	1	1	1	1	1	0	0	0	
3	0	0	0	0	1	1	1	1	1	1	1	1	5	5	5	
2	0	0	0	0	0	0	0	0	0	0	0	0	5	4	4	
1	0	0	0	0	0	0	0	0	0	0	0	0	5	4	3	
0	0	0	0	0	0	0	0	0	0	0	0	0	5	4	2	
	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15

CS 336/436, GD Hager with slides from Ji Yeong Lee, Howie Choset and Z. Dodds

The Wavefront in Action (Part 4)

- And again...

7	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
6	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
5	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
4	0	0	0	0	1	1	1	1	1	1	1	1	6	6	6	
3	0	0	0	0	1	1	1	1	1	1	1	1	5	5	5	
2	0	0	0	0	0	0	0	0	0	0	0	0	6	5	4	
1	0	0	0	0	0	0	0	0	0	0	0	0	6	5	4	
0	0	0	0	0	0	0	0	0	0	0	0	0	6	5	2	
	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15

CS 336/436, GD Hager with slides from Ji Yeong Lee, Howie Choset and Z. Dodds

The Wavefront in Action (Part 5)

- And again until...

7	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
6	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
5	0	0	0	0	0	0	0	0	0	0	7	7	7	7	7	
4	0	0	0	0	1	1	1	1	1	1	1	1	1	6	6	
3	0	0	0	0	1	1	1	1	1	1	1	1	5	5	5	
2	0	0	0	0	0	0	0	0	0	0	7	6	5	4	4	
1	0	0	0	0	0	0	0	0	0	0	7	6	5	4	3	
0	0	0	0	0	0	0	0	0	0	0	7	6	5	4	2	
	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15

CS 336/436, GD Hager with slides from Ji Yeong Lee, Howie Choset and Z. Dodds

The Wavefront in Action (Done)

- You're done
 - Remember, 0's should only remain if unreachable regions exist


7	18	17	16	15	14	13	12	11	10	9	9	9	9	9	9	
6	17	17	16	15	14	13	12	11	10	9	8	8	8	8	8	
5	17	16	16	15	14	13	12	11	10	9	8	7	7	7	7	
4	17	16	15	15	1	1	1	1	1	1	1	1	6	6	6	
3	17	16	15	14	1	1	1	1	1	1	1	1	5	5	5	
2	17	16	15	14	13	12	11	10	9	8	7	6	5	4	4	
1	17	16	15	14	13	12	11	10	9	8	7	6	5	4	3	
0	17	16	15	14	13	12	11	10	9	8	7	6	5	4	2	
	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15

CS 336/436, GD Hager with slides from Ji Yeong Lee, Howie Choset and Z. Dodds

The Wavefront, Now What?


- To find the shortest path, according to your metric, simply always move toward a cell with a lower number
 - The numbers generated by the Wavefront planner are roughly proportional to their distance from the goal

Two possible shortest paths shown



CS 336/436, GD Hager with slides from Ji Yeong Lee, Howie Choset and Z. Dodds

Another Example



Wavefront (Overview)

- Divide the space into a grid.
- Number the squares starting at the start in either 4 or 8 point connectivity starting at the goal, increasing till you reach the start.
- Your path is defined by any uninterrupted sequence of decreasing numbers that lead to the goal.

CS 336/436, GD Hager with slides from Ji Yeong Lee, Howie Choset and Z. Dodds

Navigation Functions

- A function $\phi: Q_{\text{free}} \rightarrow [0, 1]$ is called a *navigation function* if it
 - is smooth (or at least C^2)
 - has a unique minimum at q_{goal}
 - is uniformly maximal on the boundary of free space
 - is Morse
- A function is Morse if every critical point (a point where the gradient is zero) is isolated.
- The question: when can we construct such a function?

CS 336/436, GD Hager with slides from Ji Yeong Lee, Howie Choset and Z. Dodds

Sphere World

- Suppose that the world is a sphere of radius r_0 centered at q_0 containing n obstacles of radius r_i centered at $q_i, i=1 \dots n$
 - $\beta_0(q) = -d^2(q, q_0) + r_0^2$
 - $\beta_i(q) = d^2(q, q_i) - r_i^2$ $\mathcal{QO}_i = \{q \mid \beta_i(q) \leq 0\}$
- Define $\beta(q) = \prod \beta_i(q)$ (Repulsive)
 - note this is zero on any obstacle boundary, positive in free space and negative inside an obstacle
- Define $\gamma_\kappa(q) = (d(q, q_{\text{goal}}))^{2\kappa}$ (Attractive)
 - note this will be zero at the goal, and increasing as we move away
 - κ controls the rate of growth

CS 336/436, GD Hager with slides from Ji Yeong Lee, Howie Choset and Z. Dodds


Sphere World

- Consider now $\frac{\gamma_\kappa}{\beta}(q)$
 - $\frac{\gamma_\kappa}{\beta}(q)$ is only zero at the goal
 - $\frac{\gamma_\kappa}{\beta}(q)$ goes to infinity at the boundary of any obstacle
 - By increasing κ , we can make the gradient at any direction point toward the goal
 - It is possible to show that the only stationary point is the goal, with positive definite Hessian because $\partial \gamma_\kappa / \partial q$ dominates $\partial \beta / \partial q$,
 - therefore no local minima
- In short, following the gradient of $\frac{\gamma_\kappa}{\beta}(q)$ is guaranteed to get to the goal (for a large enough value of κ)

CS 336/436, GD Hager with slides from Ji Yeong Lee, Howie Choset and Z. Dodds

An Example: Sphere World

- One problem: the value $\frac{\gamma_\kappa}{\beta}(q)$ may be very large
- A solution: introduce a “switch” $\sigma_\lambda(x) = x/(\lambda + x)$



- Now, define $s(q, \lambda) = \left(\sigma_\lambda \circ \frac{\gamma_\kappa}{\beta}\right)(q) = \left(\frac{\gamma_\kappa}{\lambda\beta + \gamma_\kappa}\right)(q)$
 - this bounds the value of the function
 - however, $s(q, \lambda)$ may turn out not to be Morse

- A solution: introduce a “sharpening function” $\xi_\kappa(x) = x^{\frac{1}{\kappa}}$

$$\varphi(q) = \left(\xi_\kappa \circ \sigma_1 \circ \frac{\gamma_\kappa}{\beta}\right)(q) = \frac{d^2(q, q_{\text{goal}})}{\left[(d(q, q_{\text{goal}}))^{2\kappa} + \beta(q)\right]^{1/\kappa}}$$


For large enough κ , this is a navigation function on the sphere world!

CS 336/436, GD Hager with slides from Ji Yeong Lee, Howie Choset and Z. Dodds

Navigation Function for Sphere World


$$\varphi(q) = \left(\xi_\kappa \circ \sigma_1 \circ \frac{\gamma_\kappa}{\beta}\right)(q) = \frac{d^2(q, q_{\text{goal}})}{\left[(d(q, q_{\text{goal}}))^{2\kappa} + \beta(q)\right]^{1/\kappa}}$$

- For sufficiently large κ , $\varphi(q)$ is a navigation function




CS 336/436, GD Hager with slides from Ji Yeong Lee, Howie Choset and Z. Dodds


Navigation Function : $\varphi(q)$, varying k




k=3




k=4




k=6



k=7



k=8



k=10

CS 336/436, GD Hager with slides from Ji Yeong Lee, Howie Choset and Z. Dodds


From Spheres to Stars and Beyond

- While it may not seem like it, we have solved a very general problem
- Suppose we have a **diffeomorphism** δ from some world W to a sphere world S
 - if O''_k is a navigation function on S then
 - $O'''_k(q) = O''_k(\delta(q))$ is a navigation function on W !
 - note we also need to take the diffeomorphism into account for distances
 - Because δ is a diffeomorphism, the Jacobian is full rank
 - Because the Jacobian is full rank, the gradient map cannot have new zeros introduced (which could only happen if the gradient was in the null space of the Jacobian)
- A star world is one example where a diffeomorphism is known to exist
 - a star-shaped set is one in which all boundary points can be “seen” from some single point in the space.




$$\exists x \text{ such that } \forall y \in S, \quad tx + (1-t)y \in S \quad \forall t \in [0, 1]$$

Which of the following are the same?



CS 336/436, GD Hager with slides from Ji Yeong Lee, Howie Choset and Z. Dodds

_____jections




CS 336/436, GD Hager with slides from Ji Yeong Lee, Howie Choset and Z. Dodds

Diffeomorphism vs. Homeomorphism

HOMEOMORPHISM If $\phi: S \rightarrow T$ is a bijection, and both ϕ and ϕ^{-1} are continuous, then ϕ is a homeomorphism. When such a ϕ exists, S and T are said to be homeomorphic.


A mapping $\phi: U \rightarrow V$ is said to be *smooth* if all partial derivatives of ϕ , of all orders, are well defined (i.e., ϕ is of class C^∞). With the notion of smoothness, we define a second type of bijection.

DIFFEOMORPHISM A smooth map $\phi: U \rightarrow V$ is a diffeomorphism if ϕ is bijective and ϕ^{-1} is smooth. When such a ϕ exists, U and V are said to be diffeomorphic.



CS 336/436, GD Hager with slides from Ji Yeong Lee, Howie Choset and Z. Dodds

Which of the following are the same?



CS 336/436, GD Hager with slides from Ji Yeong Lee, Howie Choset and Z. Dodds

From Spheres to Stars and Beyond


- While it may not seem like it, we have solved a very general problem
- Suppose we have a **diffeomorphism** δ from some world W to a sphere world S
 - if O''_κ is a navigation function on S then
 - $O''_\kappa(q) = O''_\kappa(\delta(q))$ is a navigation function on W !
 - note we also need to take the diffeomorphism into account for distances
 - Because δ is a diffeomorphism, the Jacobian is full rank
 - Because the Jacobian is full rank, the gradient map cannot have new zeros introduced (which could only happen if the gradient was in the null space of the Jacobian)
- A star world is one example where a diffeomorphism is known to exist
 - a star-shaped set is one in which all boundary points can be “seen” from some single point in the space.



$$\exists x \text{ such that } \forall y \in S, \quad tx + (1-t)y \in S \quad \forall t \in [0, 1]$$

CS 336/436, GD Hager with slides from Ji Yeong Lee, Howie Choset and Z. Dodds

Construct the Mapping



CS 336/436, GD Hager with slides from Ji Yeong Lee, Howie Choset and Z. Dodds

Potential Fields on Non-Euclidean Spaces

- Thus far, we've dealt with points in \mathbb{R}^n --- what about real manipulators
- Recall we can think of the gradient vectors as forces -- the basic idea is to define forces in the workspace (which is \mathfrak{R}^2 or \mathfrak{R}^3)

force f acting at a point $x = \phi(q)$

force u acting in the robot's configuration

$\dot{x} = J\dot{q}$, where $J = \partial\phi/\partial q$

$u^T \dot{q}$ Power in configuration space

$f^T \dot{x}$ Power in workspace

Power is conserved!

$$f^T J \dot{q} = u^T \dot{q}$$

$$f^T J = u^T$$

$$J^T f = u.$$


CS 336/436, GD Hager with slides from Ji Yeong Lee, Howie Choset and Z. Dodds

Force on an Object

$$q = [x, y, \theta]^T$$

$$[a_x, a_y]^T$$

robot's local coordinate frame.



$$\phi(q) = \begin{bmatrix} x + a_x \cos \theta - a_y \sin \theta \\ y + a_x \sin \theta + a_y \cos \theta \end{bmatrix}$$

$$J(q) = \frac{\partial \phi}{\partial q}(q) = \begin{bmatrix} 1 & 0 & -a_x \sin \theta - a_y \cos \theta \\ 0 & 1 & a_x \cos \theta - a_y \sin \theta \end{bmatrix}$$

$$\begin{bmatrix} u_x \\ u_y \\ u_\theta \end{bmatrix} = J^T(q) \begin{bmatrix} f_x \\ f_y \end{bmatrix}$$

$$= \begin{bmatrix} 1 & 0 \\ 0 & 1 \\ -a_x \sin \theta - a_y \cos \theta & a_x \cos \theta - a_y \sin \theta \end{bmatrix} \begin{bmatrix} f_x \\ f_y \end{bmatrix}$$

$$= \begin{bmatrix} f_x \\ f_y \\ -f_x(a_x \sin \theta + a_y \cos \theta) + f_y(a_x \cos \theta - a_y \sin \theta) \end{bmatrix}$$


torque

CS 336/436, GD Hager with slides from Ji Yeong Lee, Howie Choset and Z. Dodds

Potential Fields for Multiple Bodies

- Recall we can think of the gradient vectors as forces -- the basic idea is to define forces in the workspace (which is \mathbb{R}^2 or \mathbb{R}^3)
 - We have $J^t f = u$ where f is in W and u is in Q
 - Thus, we can define forces in W and then map them to Q
 - Example: our two-link manipulator

$$\begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} L_1 c_\alpha \\ L_1 s_\alpha \end{bmatrix} + \begin{bmatrix} L_2 c_{\alpha+\beta} \\ L_2 s_{\alpha+\beta} \end{bmatrix}$$



CS 336/436, GD Hager with slides from Ji Yeong Lee, Howie Choset and Z. Dodds

Potential Fields on Non-Euclidean Spaces


- Example: our two-link manipulator
- $J = \begin{bmatrix} -L_1 s_\alpha - L_2 s_{\alpha+\beta} & -L_2 s_{\alpha+\beta} \\ L_1 c_\alpha + L_2 c_{\alpha+\beta} & L_2 c_{\alpha+\beta} \end{bmatrix}$

Suppose $q_{\text{goal}} = (0,0)^t$, then $f_W = (x,y)$

$$f_q = x (-L_1 s_\alpha - L_2 s_{\alpha+\beta}) + y (L_1 c_\alpha + L_2 c_{\alpha+\beta})$$

$$x (-L_2 s_{\alpha+\beta}) + y L_2 c_{\alpha+\beta}$$

$$\begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} L_1 c_\alpha \\ L_1 s_\alpha \end{bmatrix} + \begin{bmatrix} L_2 c_{\alpha+\beta} \\ L_2 s_{\alpha+\beta} \end{bmatrix}$$




CS 336/436, GD Hager with slides from Ji Yeong Lee, Howie Choset and Z. Dodds

In General

- Pick several points on the manipulator
- Compute attractive and repulsive potentials for each
- Transform these into the configuration space and add
- Use the resulting force to move the robot (in its configuration space)

Be careful to use the correct Jacobian!




CS 336/436, GD Hager with slides from Ji Yeong Lee, Howie Choset and Z. Dodds

Potential Function on Rigid Body

pick control points $\{r_i\}$ on the robot

Pick enough points to “pin down” robot (2 in plane)



$$U_{\text{att},j}(q) = \begin{cases} \frac{1}{2}\zeta_i d^2(r_j(q), r_j(q_{\text{goal}})), & d(r_j(q), r_j(q_{\text{goal}})) \leq d_0 \\ d\zeta_i d(r_j(q), r_j(q_{\text{goal}})) - \frac{1}{2}\zeta_i d^2, & d(r_j(q), r_j(q_{\text{goal}})) > d_0. \end{cases}$$


$$\nabla U_{\text{att},j}(q) = \begin{cases} \zeta_i (r_j(q) - r_j(q_{\text{goal}})), & d(r_j(q), r_j(q_{\text{goal}})) \leq d_0, \\ \frac{d\zeta_j (r_j(q) - r_j(q_{\text{goal}}))}{d(r_j(q), r_j(q_{\text{goal}}))}, & d(r_j(q), r_j(q_{\text{goal}})) > d_0. \end{cases}$$

$$U_{\text{repi},j}(q) = \begin{cases} \frac{1}{2}\eta_j \left(\frac{1}{d_i(r_j(q))} - \frac{1}{Q_i^*} \right)^2, & d_i(r_j(q)) \leq Q_i^* \\ 0, & d_i(r_j(q)) > Q_i^* \end{cases}$$

$$\nabla U_{\text{repi},j}(q) = \begin{cases} \eta_j \left(\frac{1}{Q_i^*} - \frac{1}{d_i(r_j(q))} \right) \frac{1}{d_i^2(r_j(q))} \nabla d_i(r_j(q)), & d_i(r_j(q)) \leq Q_i^*, \\ 0, & d_i(r_j(q)) > Q_i^*. \end{cases}$$

$$u(q) = \sum_i u_{\text{atti}}(q) + \sum_j u_{\text{repi}}(q)$$

$$= \sum_i J_i^T(q) f_{\text{atti}}(q) + \sum_j J_j^T(q) f_{\text{repi}}(q)$$



More points please

A Simulation Example

- Problem: simulate a planar n-link (revolute) manipulator.
- Kinematics: Let $v(\theta) = [c_\theta, s_\theta]^t$
- Points of revolution: $p_0 = [0, 0]^t$ $\alpha_i = \sum_{j=1}^i \theta_j$
 $p_i = p_{i-1} + L_i v(\alpha_i)$
- Jacobian: $w(\theta) = [-s_\theta, c_\theta]^t$

$$J_n = L_n w(\alpha_n)$$

$$J_{n-1} = J_n + L_{n-1} w(\alpha_{n-1})$$

- Now, use the revolute points as the control points to generate force vectors (note this could lead to problems in some cases).

CS 336/436, GD Hager with slides from Ji Yeong Lee, Howie Choset and Z. Dodds

Summary

- Basic potential fields
 - attractive/repulsive forces
- Gradient following and Hessian
- Navigation functions
- Extensions to more complex manipulators

CS 336/436, GD Hager with slides from Ji Yeong Lee, Howie Choset and Z. Dodds