



# FFTs in Graphics and Vision

Invariance of Shape Descriptors



# Announcements

- Homework 2 extension:
  - Now due 04/01/19



# Outline

- Math Overview
  - Translation and Rotation Invariance
  - The 0<sup>th</sup> Order Frequency Component
- Shape Descriptors
- Invariance



# Translation Invariance

Given a function  $f$  in 2D, we obtain a translation invariant representation of the function by storing the magnitudes of the frequency components:

$$f(x, y) = \sum_{l, m=-\infty}^{\infty} \hat{\mathbf{f}}_{lm} \frac{e^{i(lx+my)}}{2\pi}$$
$$\Downarrow$$
$$\{\|\hat{\mathbf{f}}_{lm}\|\} \quad l, m \in \mathbb{Z}$$



# Rotation Invariance (Circle)

Given a function  $f(\theta)$  on a circle, we obtain a rotation invariant representation by storing the magnitudes of the frequency components:

$$f(\theta) = \sum_{l=-\infty}^{\infty} \hat{\mathbf{f}}_l \frac{e^{il\theta}}{\sqrt{2\pi}}$$

$\Downarrow$

$$\{\|\hat{\mathbf{f}}_l\|\} \quad l \in \mathbb{Z}$$




# Rotation Invariance (2D)

Given a function  $f(x, y)$  in 2D, we obtain a rotation invariant representation of  $f$  by:

- Expressing  $f$  in polar coordinates:

$$f(r, \theta) = f(r \cdot \cos \theta, r \cdot \sin \theta)$$





# Rotation Invariance (2D)

Given a function  $f(x, y)$  in 2D, we obtain a rotation invariant representation of  $f$  by:

- Expressing  $f$  in polar coordinates:

$$f(r, \theta) = f(r \cdot \cos \theta, r \cdot \sin \theta)$$

- Expressing each radial restriction in terms of its Fourier decomposition:

$$f(r, \theta) = \sum_{l=-\infty}^{\infty} \hat{f}_l(r) \frac{e^{il\theta}}{\sqrt{2\pi}}$$

- Storing the magnitude of the frequency components of the different radial restrictions:

$$\left\{ \|\hat{f}_l(r)\| \cdot \sqrt{2\pi r} \right\} \quad l \in \mathbb{Z}, r \in [0, 1]$$



# Rotation Invariance (Sphere)

Given a function  $f(\theta, \phi)$  on a sphere, we obtain a rotation invariant representation by storing the magnitudes of the frequency components:

$$f(\theta, \phi) = \sum_{l=0}^{\infty} \sum_{m=-l}^l \hat{\mathbf{f}}_{lm} \cdot Y_l^m(\theta, \phi)$$

$\Downarrow$

$$\left\{ \sqrt{\sum_{m=-l}^l \|\hat{\mathbf{f}}_{lm}\|^2} \right\} \quad l \in \mathbb{Z}^{\geq 0}$$






# Rotation Invariance (3D)

Given a function  $f(x, y, z)$  in 3D, we obtain a rotation invariant representation of  $f$  by:

- Expressing  $f$  in spherical coordinates:

$$f(r, \theta, \phi) = f(r \cdot \cos \theta \cdot \sin \phi, r \cdot \cos \phi, r \cdot \sin \theta \cdot \sin \phi)$$





# Rotation Invariance (3D)

Given a function  $f(x, y, z)$  in 3D, we obtain a rotation invariant representation of  $f$  by:

- Expressing  $f$  in spherical coordinates:  
 $f(r, \theta, \phi) = f(r \cdot \cos \theta \cdot \sin \phi, r \cdot \cos \phi, r \cdot \sin \theta)$
- Expressing each radial restriction in terms of its spherical harmonic decomposition:

$$f(r, \theta, \phi) = \sum_{l=0}^{\infty} \sum_{m=-l}^l \hat{\mathbf{f}}_{lm}(r) \cdot Y_l^m(\theta, \phi)$$

- Storing the size of the frequency components coefficients of the different radial restrictions:

$$\left\{ \sqrt{\sum_{m=-l}^l \|\hat{\mathbf{f}}_{lm}(r)\|^2} \cdot \sqrt{4\pi r^2} \right\} \quad l \in \mathbb{Z}^{\geq 0}, r \in [0,1]$$



# The 0<sup>th</sup> Order Frequency Component

Given a function on the circle  $f(\theta)$ , we can express the function in terms of its Fourier decomposition:

$$f(\theta) = \sum_{l=-\infty}^{\infty} \hat{f}_l \frac{e^{il\theta}}{\sqrt{2\pi}}$$

What is the meaning of the 0<sup>th</sup> order frequency component?



# The 0<sup>th</sup> Order Frequency Component

The  $l^{\text{th}}$  frequency is the dot product of the function with the  $l^{\text{th}}$  complex exponential:

$$\hat{\mathbf{f}}_l = \left\langle f(\theta), \frac{e^{il\theta}}{\sqrt{2\pi}} \right\rangle = \int_0^{2\pi} f(\theta) \cdot \frac{e^{-il\theta}}{\sqrt{2\pi}} d\theta$$

So the 0<sup>th</sup> frequency component is:

$$\hat{\mathbf{f}}_0 = \frac{1}{\sqrt{2\pi}} \int_0^{2\pi} f(\theta) d\theta$$

# The 0<sup>th</sup> Order Frequency Component



Up to a normalization term, the 0<sup>th</sup> frequency component of a function  $f(\theta)$  is the integral of the function over the circle:

$$\hat{\mathbf{f}}_0 = \frac{1}{\sqrt{2\pi}} \int_0^{2\pi} f(\theta) d\theta$$

# The 0<sup>th</sup> Order Frequency Component



Given a function on the sphere  $f(\theta, \phi)$ , we can express the function in terms of its spherical harmonic decomposition:

$$f(\theta, \phi) = \sum_{l=0}^{\infty} \sum_{m=-l}^l \hat{\mathbf{f}}_{lm} \cdot Y_l^m(\theta, \phi)$$

What is the meaning of the 0<sup>th</sup> order frequency component?



# The 0<sup>th</sup> Order Frequency Component

The  $(l, m)^{\text{th}}$  frequency component is computed by taking the dot product of the function with the  $(l, m)^{\text{th}}$  spherical harmonic:

$$\hat{\mathbf{f}}_{lm} = \langle f(\theta, \phi), Y_l^m(\theta, \phi) \rangle$$

So the 0<sup>th</sup> frequency component is:

$$\hat{\mathbf{f}}_{00} = \frac{1}{\sqrt{4\pi}} \int_{|p|=1} f(p) dp$$

# The 0<sup>th</sup> Order Frequency Component



Up to a normalization term, the 0<sup>th</sup> frequency component of a function  $f(\theta, \phi)$  is the integral of the function over the sphere:

$$\hat{\mathbf{f}}_{00} = \frac{1}{\sqrt{4\pi}} \int_{|p|=1} f(p) dp$$



# The 0<sup>th</sup> Order Frequency Component



Note:

In the case that the function  $f$  is positive the 0<sup>th</sup> frequency coefficient will also be positive:

$$\begin{aligned}\|\hat{\mathbf{f}}_0\| &= \hat{\mathbf{f}}_0 \\ \|\hat{\mathbf{f}}_{00}\| &= \hat{\mathbf{f}}_{00}\end{aligned}$$



# Outline

- Math Overview
- Shape Descriptors
  - Shape Histograms (Ankerst *et al.*)
  - Shape Distributions (Osada *et al.*)
  - Extended Gaussian Images (Horn)
- Invariance



# Shape Matching

## General Approach

Define a function that takes in two models and returns a measure of their proximity.

$$D\left(\begin{array}{|c|} \hline \begin{array}{c} \text{Car Model } M_1 \end{array} \\ \hline \end{array} \begin{array}{|c|} \hline \text{Car Model } M_2 \\ \hline \end{array}\right) \leq D\left(\begin{array}{|c|} \hline \begin{array}{c} \text{Car Model } M_1 \end{array} \\ \hline \end{array} \begin{array}{|c|} \hline \text{Dog Model } M_3 \\ \hline \end{array}\right)$$



$M_1$  is closer to  $M_2$  than it is to  $M_3$




# Shape Descriptors

## Challenge

It is difficult to match shapes directly:

- Different triangulations of the same shape
- Different shapes have different genus
- The same shape may be in different poses
- Etc.







# Shape Descriptors

## Solution

Represent shapes by a structured abstraction that represents every shape in the same domain.





# Outline



- Math Overview
- Shape Descriptors
  - Shape Histograms (Ankerst *et al.*)
  - Shape Distributions (Osada *et al.*)
  - Extended Gaussian Images (Horn)
- Invariance



# Shape Histograms

## Approach

- Decompose space into concentric shells
- Store how much of the shape falls into each of the shells







# Shape Histograms

## Properties

- The shape is represented by 1D array of values.
- The representation is invariant to rotation







# Outline

- Math Overview
- Shape Descriptors
  - Shape Histograms (Ankerst *et al.*)
  - Shape Distributions (Osada *et al.*)
  - Extended Gaussian Images (Horn)
- Invariance




# D2 Shape Distributions


## Approach

Avoid the whole problem of tessellation, genus, etc. by building the shape descriptor from random samples from the surface of the model:

Triangulated Model



Point Set







# D2 Shape Distributions

## Key Idea

Use the fact that the distance between pairs of points on the model does not change if the model is translated and/or rotated.






# D2 Shape Distributions

## Descriptor

Represent shapes by the histogram of distances between pairs of points on the model:

$$D2_P(d) = \frac{|\{p, q \in P \mid \|p - q\| = d\}|}{|P|^2}$$








# D2 Shape Distributions

## Properties

- The shape is represented by 1D array of values.
- The representation is invariant to translations and rotations



Parameterization



Parameterization



# Outline

- Math Overview
- Shape Descriptors
  - Shape Histograms (Ankerst *et al.*)
  - Shape Distributions (Osada *et al.*)
  - Extended Gaussian Images (Horn)
- Invariance




# Extended Gaussian Images

## Approach

Use the fact that every point on the surface has a position and a normal.

Triangulated Model



Oriented Point Set






# Extended Gaussian Images

## Descriptor

Represent a model by binning points based on the associated surface normal








# Extended Gaussian Images

## Descriptor

Represent a model by binning points based on the associated surface normal






# Extended Gaussian Images

## Descriptor

Represent a model by binning points based on the associated surface normal






# Extended Gaussian Images

## Descriptor

Represent a model by binning points based on the associated surface normal






# Extended Gaussian Images

## Descriptor

Represent a model by binning points based on the associated surface normal






# Extended Gaussian Images

## Descriptor

Represent a model by binning points based on the associated surface normal






# Extended Gaussian Images

## Properties

- A 2D curve / 3D surface is represented by a histogram over a circle / sphere.
- The representation is invariant to translations.





# Outline

- Math Overview
- Shape Descriptors
- **Invariance**



# Normalization vs. Invariance

We say that a shape representation is normalized with respect to translation / rotation if the shape is placed into a canonical pose.






# Normalization vs. Invariance

We say that a shape representation is normalized with respect to translation / rotation if the shape is placed into a canonical pose.

Example:

We can normalize for translation by moving the surface so that the center of mass is at the origin.





# Normalization vs. Invariance

We say that a shape representation is invariant with respect to translation / rotation if the representation discards information that depends on translation / rotation.



# Invariance


We have seen a general method for making functions invariant to translation and rotation.



# Invariance

## Translation:

Compute the Fourier decomposition and store just the magnitudes of the Fourier coefficients.



Cartesian Coordinates

$$f(x, y, z) = \sum_{l,m,n} \hat{\mathbf{f}}_{lmn} \cdot \frac{e^{i(lx+my+zn)}}{(2\pi)^{1.5}}$$

$$\{\|\hat{\mathbf{f}}_{lmn}\|\}_{l,m,n}$$


Translation Invariant  
Representation



# Invariance

## Rotation:

Compute the spherical harmonic decomposition and store just the sizes of the different frequency components of the different radial restrictions.



Spherical Coordinates

$$f(r, \theta, \phi) = \sum_{l=0}^{\infty} \sum_{m=-l}^m \hat{\mathbf{f}}_{lm}(r) \cdot Y_l^m(\theta, \phi)$$

$$\left\{ \sqrt{\sum_{m=-l}^l \|\hat{\mathbf{f}}_{lm}(r)\|^2 \cdot \sqrt{4\pi r^2}} \right\}_{l=0}^{\infty}$$

Rotation Invariant Representation



# Overblown Claim

All methods that represent 3D shapes in either a translation-invariant or rotation-invariant method implicitly use these invariance approaches.



# Goal

Given the three shape descriptors:



- Shape Histograms
  - Shape Distributions
  - Extended Gaussian Images
- How does the descriptor obtain its invariance?
  - How can the descriptiveness of the descriptor be improved while maintaining invariance?



# Shape Histograms

The shape descriptor represents a 3D shape by binning points by their distance from the center.

It is rotation invariant.









# Shape Histograms

The shape histogram starts by representing the surface by a 3D function, obtained by rasterizing the boundary into a voxel grid:

- A voxel has value 1 if intersects the boundary
- A voxel has value 0 otherwise.



Model



Rasterization



# Shape Histograms

The shape histogram can be obtained by setting the value of the bin corresponding to radius  $r$  equal to the “size” of the rasterization restricted to the sphere of radius  $r$ :

$$\text{ShapeHistogram}(r) = \int_{|p|=r} \text{Raster}(p) dp$$



# Shape Histograms

We can express the rasterization in spherical coordinates:

$$R(r, \theta, \phi) = \text{Raster}(r \cdot \cos \theta \cdot \sin \phi, r \cdot \cos \phi, r \cdot \sin \theta \cdot \sin \phi)$$

Then, fixing the radius, we can express the function in terms of spherical harmonics:

$$R(r, \theta, \phi) = \sum_{l=0}^{\infty} \sum_{m=-l}^l \hat{\mathbf{R}}_{lm}(r) \cdot Y_l^m(\theta, \phi)$$



# Shape Histograms

In this formulation, the value of the shape histogram at a radius of  $r$  is the value of the 0<sup>th</sup> spherical harmonic coefficient:\*

$$\text{ShapeHistogram}(r) = \hat{\mathbf{R}}_{00}(r) \cdot \sqrt{4\pi r^2}$$

\*The scale factor of  $\sqrt{4\pi r^2}$  accounts for the fact that the area of the sphere of radius  $r$  is  $4\pi r^2$ .



# Shape Histograms

So the shape histogram obtains its rotation invariance by storing the (size of the) 0<sup>th</sup> order frequency component:

$$\text{ShapeHistogram}(r) = \hat{\mathbf{R}}_{00}(r) \cdot \sqrt{4\pi r^2}$$

## Extension:

We can obtain a more descriptive representation, without giving up rotation invariance, by storing the size of every frequency component:


$$\text{EShapeHistogram}(r, l) = \sqrt{\sum_{m=-l}^l \|\hat{\mathbf{R}}_{lm}(r)\|^2} \cdot \sqrt{4\pi r^2}$$




# D2 Shape Distribution

The shape descriptor represents a 3D shape by binning point-pairs by their distance.

It is both translation and rotation invariant.



3D Model



D2 Distribution

# *D2* Shape Distribution




Let's consider the rotation invariance first.




# *D2* Shape Distribution

We can think of the *D2* shape descriptor by binning the difference vector between pairs of points on the surface.



3D Model



Binned Difference Vectors







# *D2* Shape Distribution

One way to think of the *D2* shape descriptor is by binning the difference vector between pairs of points on the surface.


Then the shape distribution can be obtained by computing the Shape Histogram of the binning:



3D Model



Binned Difference Vectors





## *D2* Shape Distribution

As with the Shape Histogram, the *D2* Shape Distribution can be realized by storing 0<sup>th</sup> order frequency components of the spherical harmonic decomposition.

### Extension:

As with the Shape Histogram the representation can be made more descriptive, without sacrificing rotation invariance, by storing the size of every frequency component.

# *D2* Shape Distribution




What about the translation invariance?




## D2 Shape Distribution

The Shape Distribution is computed from the binning of point-pair differences. How is this function computed?



3D Model




Binned Difference Vectors




# D2 Shape Distribution

A point  $q$  on the surface will contribute to bin  $v$  if the point  $q - v$  is also on the surface.



3D Model




Binned Difference Vectors




# D2 Shape Distribution

Consider the rasterization of the surface into a regular voxel grid.



Model



Rasterization



## D2 Shape Distribution


A point  $q$  on the surface will contribute to bin  $v$  if the point  $q - v$  is also on the surface.




$$\text{Raster}(q - v) = 1$$



$$\text{DBin}(v) = \int_{q \in \text{Surface}} \text{Raster}(q - v) dq$$



3D Model



Binned Difference Vectors



# D2 Shape Distribution


For a point  $q \in \mathbb{R}^3$ , the point will only contribute to bin  $v$  if  $q$  and  $q - v$  are both on the surface.

That, is  $q$  contribute to bin  $v$  if and only if:


$$\text{Raster}(q) \cdot \text{Raster}(q - v) = 1$$



$$\text{DBin}(v) = \int_{q \in \mathbb{R}^3} \text{Raster}(q) \cdot \text{Raster}(q - v) dq$$



3D Model



Binned Difference Vectors





## D2 Shape Distribution

Thus the binning function is the correlation of the rasterization with itself:

$$\begin{aligned}\text{DBin}(v) &= \int_{q \in \mathbb{R}^3} \text{Raster}(q) \cdot \text{Raster}(q - v) \, dq \\ &= (\text{Raster} \star \text{Raster})(v)\end{aligned}$$



# D2 Shape Distribution

Recall:

To compute the correlation of  $f$  with  $g$  we multiply the Fourier coefficients of  $f$  by the conjugates of the Fourier coefficients of  $g$ :

$$(f \star g)(\theta) = \sum_{l=-\infty}^{\infty} \sqrt{2\pi} (\hat{\mathbf{f}}_l \cdot \bar{\hat{\mathbf{g}}}_l) e^{il\theta}$$

When  $f = g$ , this gives:

$$(f \star g)(\theta) = \sum_{l=-\infty}^{\infty} \sqrt{2\pi} \|\hat{\mathbf{f}}_l\|^2 e^{il\theta}$$



## D2 Shape Distribution

⇒ The binning function implicitly converts the rasterization function into a function whose Fourier coefficients are the square norms of the Fourier coefficients of the rasterization.


Which is what we do to make a function translation invariant.



# Extended Gaussian Image

This spherical shape descriptor represents a 3D shape by a histogram on the sphere.

It is obtained by binning points by their normal direction, and is translation invariant.






# Extended Gaussian Image

To obtain the EGI representation, we can think of points on the model as living in a 5D space:

- The first 3 dimensions are indexed by the position.
- The last 2 are indexed by the normal direction.






# Extended Gaussian Image

To obtain the EGI representation, we can think of points on the model as living in a 5D space.

If we fix the normal angle, we get a 3D slice of the 5D space, corresponding to all the points on the surface with the same normal:






# Extended Gaussian Image

For each normal  $n$ , the EGI stores the “size” of the points in the normal slice corresponding to  $n$ .

This is the 0<sup>th</sup> order frequency component of the rasterization of the points on the model with normal  $n$ .





# Extended Gaussian Image

For each normal  $n$ , the EGI stores the “size” of the points in the normal slice corresponding to  $n$ .

This is the 0<sup>th</sup> order frequency component of the rasterization of the points on the model with normal  $n$ .

## Extension:

We can get a more discriminating descriptor, without giving up translation invariance, by storing the size of every frequency component.