

Notation: -  $d_G(u)$  = degree of  $u$  in  $G$

$$- \Delta(G) = \max_{u \in V} d_G(u) = \max \text{ degree}$$


### Minimum Degree Spanning Tree:

Input:  $G = (V, E)$

Feasible solution: Spanning tree  $T$  of  $G$

Objective:  $\min \Delta(T)$

NP-hard (see homework)



Def: For spanning tree  $T$ , non-tree edge  $e$ , the **fundamental cycle** of  $e$  is the unique cycle in  $T \cup \{e\}$

### Local move:

pair  $(e, e')$ ,  $e \in T$ ,  $e'$  in fundamental cycle of  $e$ .

Add  $e$ , remove  $e'$


## Local Search algorithm :

Init  $T$  an arbitrary spanning tree

while there is a local move which decreases  $\Delta(T)$


do it

Problem: might be bad approximation!




Local opt: max degree  $d = \Theta(\sqrt{n})$

Global opt: max degree 3



Def: Let  $u \in V$ .  $(u, \{v, w\})$  is a  **$u$ -improvement** if  $\exists \{u, x\}$  on fundamental cycle of  $\{v, w\}$ , and if we add  $\{v, w\}$  and remove  $\{u, x\}$  to get  $T'$  then  $\max(d_{T'}(v), d_{T'}(w)) \leq d_{T'}(u) = d_T(u) - 1$

Example:



LS Alg 2:

while  $\exists u$ -improvement for some node  $u$ , do it

Problem: running time!

LS Alg }:

while  $\exists$   $u$ -improvement where  $d_T(u) \geq \Delta(T) - \log n$   
do it.

Thm: Running time is polynomial

Pf:

Each iteration poly time  $\Rightarrow$  just need to bound # iterations

Potential function:

$$\Phi(u) = 3^{d_T(u)}$$

$$\Phi(T) = \sum_{u \in V} \Phi(u) = \sum_{u \in V} 3^{d_T(u)}$$

Easy facts:

$$- \Phi(T) \geq 3n$$

$$- \Phi(T) \leq n \cdot 3^n$$

Claim: Suppose did a  $u$ -improvement on  $T$  to get  $T'$ ,  
where  $d_T(u) \geq \Delta(T) - \log n$ . Then

$$\Phi(T') \leq \left(1 - \frac{2}{9n^3}\right) \Phi(T)$$

Pf of claim:

Let  $(u, \{v, w\})$  be the  $u$ -improvement, and

$$\text{let } i = d_T(u) \geq \Delta(T) - \log n$$

Decrease in  $\Phi$  ( $\Phi(T) - \Phi(T')$ )

$$u: \Phi_T(u) - \Phi_{T'}(u) = 3^i - 3^{i-1} = 2 \cdot 3^{i-1}$$

$$x: \geq 0$$

$$v: 3^{d_T(v)} - 3^{d_{T'}(v)} = \left( 3^{d_{T'}(v)} - 3^{d_T(v)} \right)$$

$$= - \left( 3^{d_T(v)} - 3^{d_{T'}(v)-1} \right)$$

$$= - 3^{d_{T'}(v)-1} (3 - 1)$$

$$= -2 \cdot 3^{d_{T'}(v)-1}$$

$$\geq -2 \cdot 3^{d_T(v)-1}$$

$$= -2 \cdot 3^{i-2}$$

$w$ : same

All other nodes: unchanged

$$\begin{aligned} \Rightarrow \Phi(T) - \Phi(T') &\geq 2 \cdot 3^{i-1} - 4 \cdot 3^{i-2} \\ &= \frac{2}{3} \cdot 3^i - \frac{4}{9} \cdot 3^i \\ &= \frac{2}{9} \cdot 3^i \end{aligned}$$

$$\geq \frac{2}{9} \cdot 3^{D(T) - \log n}$$

$$= \frac{2}{9 \cdot 3^{\log n}} \cdot 3^{D(T)}$$

$$= \frac{2}{9 \cdot n^{\log 3}} \cdot 3^{D(T)}$$

$$\geq \frac{2}{9 n^2} \cdot \frac{1}{n} \mathbb{I}(T)$$

$$= \frac{2}{9 n^3} \mathbb{I}(T)$$

$$\Rightarrow \mathbb{I}(T) \leq \left(1 - \frac{2}{9 n^3}\right) \mathbb{I}(T)$$

✓

Use claim: Suppose runs for  $\frac{9}{2} n^4 \ln 3$  iterations

$$\Rightarrow \mathbb{I}(T) \leq \left(1 - \frac{2}{9 n^3}\right)^{\frac{9}{2} n^4 \ln 3} \cdot n \cdot 3^n$$

$$\leq e^{-n \ln 3} \cdot n \cdot 3^n$$

$$= n$$


Since  $\mathbb{I}(T) \geq 3n$ , must have already finished ✓

Approximation:

Let  $T$  be tree returned by alg,

$T^*$  optimal tree,  $\Delta^* = \Delta(T^*)$

Then:  $\Delta(T) \leq 2 \cdot \Delta^* + \log n$




Lemma: Let  $(V_1, V_2, \dots, V_k)$  be a partition of  $V$ ,

let  $E' \subseteq E$  edges across partitions,

let  $V'$  a vertex cover of  $E'$ .

Then  $\Delta^* \geq \frac{k-1}{|V'|}$

Pf:



$T^*$  has  $\geq k-1$  edges of  $E'$

Each of which has  $\geq 1$   
endpoint in  $V'$

$\Rightarrow$  some vertex  $v \in V'$  has

$$d_{T^*}(v) \geq \frac{k-1}{|V'|}$$

From now on:  $i \geq \Delta(T) - \log n$

Def: Let  $S_i = \{v \in V : d_T(v) \geq i\}$

Def: Let  $E_i^T$  be **edges of  $T$**  incident on at least one node in  $S_i$

Claim:  $|E_i^T| \geq (i-1)|S_i| + 1$


Pr:  $\sum_{v \in S_i} d_T(v) \geq i|S_i|$  (**def of  $S_i$** )

$|\{u,v \in E_i^T : u,v \in S_i\}| \leq |S_i| - 1$  ( **$T$  a tree**)

$$\Rightarrow |E_i^T| = \sum_{v \in S_i} d_T(v) - |\{u,v \in E_i^T : u,v \in S_i\}|$$

$$\geq i|S_i| - (|S_i| - 1) = (i-1)|S_i| + 1$$

Def: Let  $E_i^G$  be **edges in  $G$**  between components of  $T - E_i^T$



(partition we'll  
apply first  
lemma to)



Claim:  $S_{i-1}$  is a vertex cover of  $E_i^G$

Pr: Let  $e = \{x, y\} \in E_i^G$

Case 1:  $e \in E_i^T$ .

By def of  $E_i^T$ , either  $x$  or  $y$  (or both)  
is in  $S_i \Rightarrow$  in  $S_{i-1}$  ✓

Case 2:  $e \notin E_i^T$

Since  $e \notin T$ , has fundamental cycle  $C$

$\Rightarrow$  since  $x, y$  in diff components of  $T - E_i^T$ ,

$C$  has some edge  $\{u, v\}$  from  $E_i^T$

$\Rightarrow$  either  $u$  or  $v$  in  $S_i$  : w.l.o.g,  $u \in S_i$

$\Rightarrow$  since at local opt,  $(u, \{x, y\})$  cannot  
be  $u$ -improvement

$\Rightarrow \max(d_T(x)+1, d_T(y)+1) \geq d_T(u) \geq i$

$\Rightarrow$  either  $x$  or  $y$  in  $S_{i-1}$

Claim:  $\exists i \geq \Delta(T) - \log n$  s.t.  $|S_{i-1}| \leq 2|S_i|$

Pf:

$S_p$  false  $\Rightarrow |S_{i-1}| > 2|S_i| \quad \forall i$

$$|S_{\Delta(T)}| \geq 1 \Rightarrow |S_{\Delta(T) - \log n}| > 2^{\log n} \cdot 1 = n$$

$\Rightarrow \Leftarrow$

Can finally apply partition lemma!

Let  $i^* \geq \Delta(T) - \log n$  s.t.  $|S_{i^*-1}| \leq 2|S_{i^*}|$

Partition  $V$  into  $|E_{i^*}^T| + 1$  sets by  $T - E_{i^*}^T$

$S_{i^*-1}$  a vertex cover of  $E_{i^*}^G$

$$\Rightarrow \Delta^* \geq \frac{|E_{i^*}^T|}{|S_{i^*-1}|}$$

$$\geq \frac{(i^* - 1)|S_{i^*}| + 1}{2|S_{i^*}|}$$

$$\geq \frac{i^* - 1}{2} + \frac{1}{2|S_{i^*}|}$$

$$\geq \frac{\Delta(T) - \log n - 1}{2}$$

$$\Rightarrow \Delta(T) \leq 2\Delta^* + \log n + 1$$

~~2~~

$$\text{But for: } \Delta(T) \leq \Delta^* + 1$$

- Min-degree Steiner Tree

$$\text{Rightly noted (D-H-I-N)}^{(19)}; \Delta(T) \leq O(\Delta^* + \log n)$$